

**Explanation of Units**

**Intrusive Rocks**

**Devonian(?)**  
**Dg** **Two-mica granite.** Typically with feldspar phenocrysts up to 3 cm long, and coarse pegmatite. Commonly foliated, phenocrysts and foliation form a swirly pattern rather than a preferred orientation. Pegmatite is common in the stratified units, especially in the Rindgemere Formation, as migmatite or isolated bodies.  
 The best places to observe the granite are along Milton Mills Road ½ mile northwest of Nisbit Pond, and in the pavement outcrop on Sam Wentworth Road 1 ½ mile north-northwest of West Lebanon.

**Stratified Rocks**

**Devonian(?)**  
**DI** **Littleton Formation.** Gray-weathering quartz-muscovite-biotite ± garnet + staurolite + sillimanite schist, with sparse light gray to white micaceous quartzite beds. Graded beds are common. Equivalent to upper part of the Rindgemere Formation of Katz (1917) and Hussey (1962, 1987). Exposed along Key Brook in the southeast corner of the quadrangle. In the next quadrangle to the east, tops toward the southeast provide one basis for determining the topping direction of the stratigraphic sequence.

**Devonian-Silurian(?)**  
**Dsm** **Madrid Formation.** Well bedded, pale gray calc-silicate granofels and darker gray biotite-quartz-feldspar granofels. Exposed along Key Brook in the southeast corner of the quadrangle.

**Silurian(?)**  
**St** **Towow Formation.** Rusty-weathering phyllite or fine-grained schist dominated by muscovite, quartz, pyrite and/or pyrrhotite, locally with graphite, biotite, and aluminous index minerals. Contains rusty-weathering beds of sulfidic quartzite and granofels in some places.  
 The Towow Formation is best exposed in brooks, particularly Bog Brook down from Depot Road in North Lebanon and in Little River parallel to Goding Road near the Acton-Lebanon town line. Outcrops along Goding Road are also recommended. Several contractors and homeowners report high iron content in water wells drilled in the Towow Formation.

**Stg** **Grit member.** Rusty to white-weathering, sulfidic quartzites with granule-sized clasts of quartz, quartzite, and rare phyllite chips. Some quartz clasts are bluish. Grits of the Towow Formation are best exposed along Merchants Row, ¼ mile southwest of Center Lebanon. Because Towow phyllites are exposed both west and east of the grits, the grits are interpreted as a layer within the Towow Formation. A possible alternative is that they represent a horizon at the contact with the Littleton Formation, preserved in a downward-facing antiform.

**Sgo** **Gully Oven formation** (new name, informal). Well-bedded, gray to somewhat rusty-weathering schist and gray to white quartzite beds with sharp bedding contacts, graded beds not convincing anywhere in Milton quadrangle. Schist typically contains andalusite, or sillimanite after andalusite, or pseudomorphic lumps composed of muscovite, quartz, etc. Pink cotitic lenses are quite common locally; biotitic granofels and calc-silicate rocks are more rare. Colored dots on the map indicate cotitic localities. Lower contact, with the Rindgemere Formation, is gradational.  
 The Gully Oven formation is well exposed at the type locality, a deep bedrock ravine along a branch of Great Brook, west of Gully Oven Road, 2 miles north of West Lebanon (Stop 9 of Gilman, 1986). A more complete exposure across strike is afforded by the next branch of Great Brook to the northeast of Gully Oven. Another good place to see this unit is in a big ledge behind the Burke house west of Goding Road, just south of the Lebanon-Acton town line. The Burkes report black manganese stains on plumbing fixtures, presumably from well water in this unit.

**Sr** **Rindgemere Formation.** Variable unit dominated by massive, coarse-grained gray or rusty feldspathic sillimanite schist, locally with biotitic granofels and quartzite beds, and calc-silicate beds and lenses. Rustiness tends to be reddish. Rindgemere schist and granofels can be observed in the south part of Acton in a pavement outcrop near the junction of Lebanon Road and Blueberry Hill, and along Blueberry Hill Road ½ mile northeast of Black Pond. Schist containing dark gray calc-silicate lenses is well exposed in a ledge south of County Road 1 mile west of Hurd Hill. Coarse-grained, migmatitic Rindgemere Formation can be seen along the Salmon Falls River below the dam ½ mile south of Milton.

**Explanation of Lines**

**Intrusive or stratigraphic contact.** Solid where well constrained by outcrops or dense float; dashed where less well constrained, but projected with confidence; dotted where poorly constrained.

**Fault.** Solid where well constrained by silicified material or unit discontinuities; dashed where conjectural. **U**=upthrow side, **D**=downthrow side.

**Anticline axial trace.** Arrow head indicates direction of plunge.

**Structural dome in foliation.**

**Lithologic Symbols**

● **Outcrop of mapped unit.** Dot = single outcrop. Shaded area = many outcrops.

○ **Outcrop of Gully Oven formation containing cotilic.**

○ **Site of bedrock drill core** (Sevee and Maher Engineers, 1986).

≡ **Area of float presumed to represent underlying bedrock.**

⊕ **Pegmatite too small to map.** Presumably related to granite (Dg). (Vertical dike, isolated outcrop)

⚡ **Quartz vein (± also contains abundant black tourmaline).** Strike and dip indicated. (Inclined, Vertical, Dip unknown)

⚡ **Basalt dike too small to map.** Presumed to be Mesozoic. Dikes are common 1 to 1½ mile south of Hurd Hill, in outcrops along the Little River and along Winchell Orchard Road, Acton. Subsurface dikes at the proposed Hebo-Hybo Road Landfill site, Lebanon, were detected by geophysical surveys and drilling, although one does occur in outcrop (Sevee and Maher Engineering, 1986). Strike and dip indicated. (Inclined, Vertical)

⚡ **Fine-grained dioritic dike.** One dike too small to map, about 1 mile south of Hurd Hill, Acton. Strike and dip indicated. (Vertical)

**Structural Symbols**

*Oriented symbols indicate strike and dip of planar features or trend and plunge of linear features.*

⚡ **Bedding, tops unknown, measured on quartzite or granofels bed.** (Inclined, Vertical)

⚡ **Bedding, tops known.** (Overturned)

⚡ **Compositional layering in schist.** Color variation interpreted as sedimentary in origin.

⚡ **Dominant foliation.** Pervasive schistosity or phyllitic foliation. (Inclined)

⚡ **Axial plane of fold with axial plane foliation.** (Dextral, Sinistral, Rotation sense unknown)

⚡ **Axial plane of fold which deforms foliation.** (Inclined, Vertical)

⚡ **Hinge of fold which deforms foliation.** (Dextral, Neutral, Rotation sense unknown)

⚡ **Crenulation cleavage.**

⚡ **Mineral lineation (q = quartz; ms = muscovite or sillimanite).** (Plunging, Horizontal)

⚡ **Boudin neck line.**

⚡ **Intersection lineation (fb = foliation-bedding; fc = foliation-cleavage).**

⚡ **Kink band.**

⚡ **Kink axis.** (Dextral, Sinistral)

⚡ **Joint or joint set.** (Inclined, Vertical)

**GEOLOGIC TIME SCALE**

| Geologic Age     | Absolute Age*        |  |
|------------------|----------------------|--|
| Cenozoic Era     | 0-65                 | *In millions of years before present. (Okulitch, A. V., 2002. Echelle des temps géologiques, 2002. Commission géologique du Canada, Dossier Public 3040) |
| Mesozoic Era     | 65-145               |  |
|                  | 145-200              |  |
| Paleozoic Era    | Triassic Period      | 200-253  |
|                  | Permian Period       | 253-300  |
|                  | Carboniferous Period | 300-360  |
|                  | Devonian Period      | 360-418  |
|                  | Silurian Period      | 418-443  |
| Precambrian time | Ordovician Period    | 443-489  |
|                  | Cambrian Period      | 489-544  |
|                  | Older than 544       |  |

**Bedrock Geology of the Milton Quadrangle, New Hampshire-Maine**

Bedrock geologic mapping by  
**Peter J. Thompson**


Digital cartography by:  
**Susan S. Tolman**

Geologic editing by:  
**Henry N. Berry IV**

Cartographic design and editing by:  
**Robert D. Tucker**

Robert G. Marvinney  
 State Geologist

Funding for the preparation of this map was provided in part by the U.S. Geological Survey STATEMAP Program, Cooperative Agreement No. 03HQAG0068.



**Maine Geological Survey**

Address: 22 State House Station, Augusta, Maine 04333  
 Telephone: 207-287-2801 E-mail: mgsg@maine.gov  
 Home page: http://www.maine.gov/doc/nrm/nrm.htm

**Open-File No. 04-77**

**2004**

**Stratigraphy**

Katz (1917) and Hussey (1962) concluded that the Towow Formation is stratigraphically above the Rindgemere Formation, and assigned rocks both northwest and southeast of the Towow to the Rindgemere. According to their interpretation, the Towow Formations within the Lebanon syncline, overturned toward the southeast. Hussey (1968, 1985) divided the Rindgemere Formation into upper and lower members, with the upper, rhythmically-bedded member on both sides of the Towow Formation. The present author, however, thinks that rocks southeast of the Towow Formation in the Milton and adjacent Sanford quadrangles are those of the Gully Oven formation, having thicker schist beds and common graded beds, and correlative instead with the Devonian Littleton Formation. Just east of the Milton quadrangle, graded beds top to the southeast, away from the Towow. If assignment to the Littleton Formation is correct, there is no symmetry across the supposed Lebanon syncline in the Milton area.  
 Because the Rindgemere/Gully Oven/Towow sequence so closely resembles the Rangely/Perry Mountain/Smalls Falls sequence, the present author prefers the correlation with units of the "Rangely sequence" of the Rochester area, as shown in the accompanying chart. The units generally dip northward and top to the southeast, meaning that the whole sequence is overturned.

**Stratigraphic Correlation Chart**

| Milton, NH/ME (this study) | Rochester, NH (Eusden, 1984) | Berwick 12 quad (Hussey, 1962; 1985) |
|----------------------------|------------------------------|--------------------------------------|
| Littleton Fm.              | Littleton Fm.                | Upper part of Rindgemere Fm.         |
| Madrid Fm.                 | Madrid Fm. (discontinuous)   | (no equivalent)                      |
| Towow Fm.                  | Smalls Falls Fm.             | Towow Fm.                            |
| Gully Oven formation       | Perry Mountain Fm.           | Upper part of Rindgemere Fm.         |
| Rindgemere Fm.             | Rangely Fm.                  | Lower Rindgemere Fm.                 |

**Structural Geology**

formation seems to occupy an open antiform plunging gently northwest. On the west flank of Hurd Hill, bedding in the Gully Oven formation dips gently west, to the northeast, in brook exposures just outside the quadrangle boundary, bedding dips northeast.

**Regional Implications**

The author spent several field days looking at the area mapped by Eusden (1984) southwest of the Milton quadrangle, and drew some what different conclusions. Using the same schist/quartzite proportion criteria applied on the Maine side of the border for distinguishing the Gully Oven and Littleton Formations, the author concluded that the inclined second-phase folds that deform the overturned sequence in Eusden's area involve the Smalls Falls and Littleton Formations rather than the Smalls Falls and Perry Mountain Formations. Without this interpretation, some significant right-lateral offset is required along a fault or faults parallel to the Salmon Falls River along the state border. The Towow Formation mapped by Eusden near Rochester lines up with Towow folded with Littleton along the Salmon Falls River. Axial traces of these folds (F?) would extend into the broad area of the Towow Formation in the Milton quadrangle, helping to explain such a broad outcrop width for what is a relatively thin unit elsewhere. Alternatively, the Towow Formation could be thicker due to a local accumulation of reducing muds during deposition.  
 The grit horizon at Merchants Row might correlate with grits recognized by Eusden near Goose Pond, New Hampshire, at what he interpreted as the Smalls Falls / Littleton contact, and by Hussey near the Salmon Falls River at the Towow / Littleton contact (Eusden and others, 1984, stop 6). If so, the grits (Stg) surrounded by the Towow Formation in the Milton quadrangle might be explained as the crest of an antiform underlain by the Littleton Formation. They are more or less on strike with a northeast-plunging antiform cored by the Littleton Formation at the Salmon Falls River.

The extent of the Silver Mine fault in either direction is unknown. Hussey (personal communication, 2004) points out that bedding in the Littleton Formation also swings around to the north and northwest as it approaches what would be the fault's extension near Lebanon, Maine (Hussey, 1962). The fault there has apparently juxtaposed the Littleton to the west against the Rindgemere to the east. Farther south, however, the Gonic/Berwick contact does not seem to be offset where the fault might

**Metamorphism**

porphyroblasts, locally replaced by sillimanite and/or muscovite, as well as garnet. The Rindgemere Formation contains abundant sillimanite and seems to have andalusite only in the most pelitic layers; it is generally more feldspathic than the Gully Oven or the Littleton Formations. The feldspar content may be responsible for the tendency toward migmatization of the Rindgemere Formation, and the abundant pegmatite and two-mica granite bodies associated with this formation may be derived quite locally.

**References Cited**

Eusden, J. D., 1984. The bedrock geology of part of the Alton, New Hampshire, and Berwick, Maine, 15' quadrangles. M. S. thesis, University of New Hampshire, Durham, 114 p., map scale 1:62,500.

Eusden, J. D., Bothner, W. A., Hussey, A. M., II, and Laird, J., 1984. Silurian and Devonian rocks in the Alton and Berwick quadrangles, New Hampshire and Maine. In: Hanson, L. S., editor, Geology of the Coastal Lowlands, Boston, MA to Kennebec, ME. Field Guide for 76th Annual Meeting, New England Intergovernmental Geological Conference, Salem State College, Salem, MA, p. 325-351.

Gilman, R. A., 1986. Bedrock geology of the Newfield and Berwick quadrangles, southern Maine. In: Newberg, D. W. (editor), Guidebook for field trips in southwestern Maine. New England Intergovernmental Geological Conference, 78th annual meeting, Bates College, Lewiston, Maine, p. 290-305.

Hall, B. A., Pollock, S. G., and Dolan, K. M., 1976. Lower Devonian Schoonook Formation and Mtatagan Sandstone, northern Maine—a flysch basin-margin delta complex. In: Page, L. R. (editor), Contributions to the stratigraphy of New England: Geological Society of America, Memoir 148, p. 57-63.

Hatch, N. L., Jr., Moech, R. H., and Lyons, J. B., 1983. Silurian-Lower Devonian stratigraphy of eastern and south-central New Hampshire: extensions from Maine. American Journal of Science, v. 283, p. 739-761.

Hussey, A. M., II, 1962. The geology of southern York County, Maine: Maine Geological Survey, Bulletin 14, 67 p., map scale 1:62,500.


Hussey, A. M., II, 1968. Stratigraphy and structure of southwestern Maine. In: Zen, E.-on, White, W. S., Hadley, J. B., and Thompson, J. B., Jr. (editors), Studies of Appalachian geology, northern and maritime. Interscience Publishers, New York, p. 291-301.

Hussey, A. M., II, 1985. The bedrock geology of the Bath and Portland 2-degree map sheets, Maine: Maine Geological Survey, Open-File Report 85-87, 82 p., map scale 1:250,000.

Hussey, A. M., II, Rand, R. J., and Austin, M. B., 1958. Maine Metal Mines and Prospects: Maine Geological Survey, Mineral Resources Index No. 3, 53 p.

Katz, F. J., 1917. Stratigraphy in southeastern New Hampshire and southwestern Maine. U. S. Geological Survey, Professional Paper 108, p. 165-177.

King, V. T., 2000. Mineralogy of Maine, Vol. 2. Mining History, Gens. and Maher Engineers, 1986. Hydrogeologic report, Hebo-Hybo Road secure landfill, Lebanon, Maine; unpublished well logs.





**Prospect and tailings still evident in Acton-Lebanon silver mine district**

Historical maps (Hussey and others, 1958; King, 2000) indicate that more mines existed in the 1800's, but they were not mapped precisely. Some mine sites visible in the 1950's have undoubtedly filled in or collapsed.

- Prospect. Trench with quartz.
- ▲ Tailings. Pile or ridge of ball quartz rubble.
- Pit. Rectangular excavation about 4' x 6', depth unknown.

**SOURCES OF INFORMATION**

Field work by P. J. Thompson (2003). Additional structural data from Eusden (1984) and subsurface lithologic information from Sevee and Maher Engineers (1986).



Topographic base from U.S. Geological Survey Milton quadrangle, scale 1:24,000 using standard U.S. Geological Survey topographic map symbols.  
 The use of industry, firm, or local government names on this map is for location purposes only and does not implicate responsibility for any present or potential effects on the natural resources.