



Environments just above the highest high water datum, but under the partial influence of marine processes and forces.

Sd	Dunes and Vegetated Beach Ridges	Unconsolidated sand or gravel deposits capping beach environments. Dunes are subject to storm waves and winds, while gravel beach ridges are subject only to storm wash. Each may be vegetated with salt-tolerant vegetation.	Sm	Fresh-Brackish Marsh	Water-saturated, organic-rich sediments characterized by broad-leaved vegetation tolerant of constant submergence in fresh water. Salinity of interstitial water is less than 5 ppt.	Se	Eolian Flat	Partially vegetated sand flats adjacent to dune fields. Subject to generally northwest winds and occasional storm flooding.
Sw	Fresh-Brackish Water	Ponded water behind beach ridges, man-made constrictions on former tidal embayments, or on marsh surfaces transitional between upland and salt marsh environments. Salinity of the water is less than 5 parts per thousand (ppt).	Sz	Man-Made Land	Structures or fill emplaced by man in the nearshore environment.	Sf	Washover Flat	Sand deposits covering salt marshes behind inlet mouths which originate from storm washover or inlet deltaic deposits on salt marshes. Subject to storm washover and spring tide flooding.
			Sx	Landslide Excavation and Deposits	Natural excavation into shoreline upland slopes created by large-scale slumping or sliding of bank material and the resulting deposits at the base of the slopes.	Sr	Fluvial Marsh	Vegetated river floodplain and bank environments, characterized by fresh-water pond vegetation such as pond

Environments between the highest high water datum and the lowest low water datum subject to twice daily tidal flooding and all other marine forces.

M1	Marsh Environments	Vegetated environments or isolate depressions within vegetated environments located above mean tide level in protected coastal areas.	B5	Low-Energy Beach	Beaches consisting of a wide variety of sediment sizes which are protected from high wave energy. Sediment characteristics are dependent upon sediment sources, which is usually from up-land scarps immediately shoreward of the beach. Low-energy beaches may exhibit growth of salt marsh grass when there is little sediment movement.	F3	Mussel Bar	Low mounds of living mussels, <i>Mytilus edulis</i> , and/or disarticulated and broken mussel shells accumulated by wave shoaling. Mussel bars generally occur at the mouths of estuaries or embayments at tidal channel margins where nutrient-laden oceanic water first flood flat environments. Mussel bars accumulate on intertidal flats.
M2	High Salt Marsh	Organic-rich sediments densely vegetated primarily with the salt marsh grass <i>Spartina patens</i> (salt-meadow grass). High salt marshes are at the same level as mean high water.	B7	Boulder Ramp	Sloping surfaces in the lower intertidal zone veneered by large boulders. This environment is seaward of gravelly or boulder beaches on high wave energy shorelines. Boulders are remnant lag deposits of eroded glacial tills. Boulder movement is limited to periods of intense storm wave activity.	F4	Channel Levee	Linear accumulations of sediment along margins of tidal channels built by several tens of centimeters above the surrounding intertidal flats. Channel levees are constructed from sediment deposited on the flat as the tide rises above the channel margins.
M3	Low Salt Marsh	Mud or muddy sand embankments sparsely to densely vegetated by the salt marsh grass <i>Spartina alterniflora</i> (salt cord-grass). Low salt marsh exists between mean tide level and mean high water.	Bw	Washover Fan	Fan-shaped deposits of gravel located behind gravel beach ridges and covering portions of marshes. Few washovers have been recognized as discrete units on sand beaches. Washover fans are deposited by storm waves. Fan sediment is derived from the beach itself.	F5	Algal Flats	High, coarse and fine-grained intertidal flats covered with the green algae, <i>Enteromorpha recta</i> .
M4	Marsh Levee	Channel-margin sediments vegetated with salt-meadow grass which exist up to several tens of centimeters above the salt marsh surface. The marsh levee consists of sandy silt- or silt-size sediment deposited from flood waters rising above channel margins, either from high river discharge into estuarine embayments or from storm-surge embayments or tides from the ocean.	Bs	Spits	Partially-submerged beach ridges which extend offshore into open water. This category includes tombolos (spits joining an island with the mainland).	F6	Veneered Ramp	Former boulder ramps presently covered by fine-grained sediment settling out of the water column.
							Miscellaneous Environment	
	Salt Pannes and Salt Ponds	Salt-water filled, non-vegetated depressions on the high salt marsh surface (salt pannes) or salt-water filled depressions anywhere in the intertidal zone (i.e. tidal pools). Salt pannes may be dry and covered with algae during late summer months.				M	Ledge	Subaerially or subaqueously exposed bedrock.
						Mc	Fluvial-Estuarine Channel	Transitional channel between river and estuarine channels. The fluvial tidal fluvial, or estuarine state de-

Md	Spillover Lobes	Lobate bars of sediment which extend from flood-tidal deltas into estuarine or tidal channel areas.
----	-----------------	---

Environments existing below lowest low water and subject to tidal current forces and wave-generated current forces.

Flat Environments		Channel Environments		Channel Slope	
	Submerged, gently sloping, or level environments composed primarily of fine sand, silt, and clay. Includes subaqueous exposures of coarse-grained, Pleistocene glacial sediments.		Linear, intertidal and subtidal depressions carrying tidal-current water.		Gently to moderately sloping wall margins of large tidal channels. Channel slopes are confined to channel wall margins composed of sediment.
Pm	Mud Flat	C1	High-Velocity Tidal Channel	Cb	Abandoned Tidal Channel
Pc	Coarse-Grained Flat	C2	Medium-Velocity Tidal Channel		Former tidal channel no longer carrying flow sufficient to erode the channel floor or margin walls. Abandoned channels usually occur in salt marsh tracts where meandering of the central drainage channel cuts off former channel segments.
Fe	Belgras Flat	C3	Low-Velocity Tidal Channel	Cf	Tidal Fluvial Channel
Fs	Seaweed Community	C4	Estuarine Channel		Lower portions of river channels under tidal influence but not carrying estuarine waters.
Fb	Upper Shoreface	C5	Estuarine Flood Channel		Tidal Creeks
	The inner subtidal slope which extends seaward from large exposed sand beaches where sediments are actively transported by bottom currents generated by storm waves. The upper shoreface is a sandy environment of constant wave shoaling under normal wave conditions.	C6	Estuarine Ebb Channel		Marsh Drainage Ditch
Fp	Lower Shoreface	C7	Inlet Channel		Approximate transition boundary between estuarine and marine (30 ppt salinity) waters and between estuarine and river (0.5 ppt) waters.
	The outer subtidal slope which extends seaward from the upper shoreface. The lower shoreface is affected only by currents generated by storm waves. Lower shoreface sediments grade from sand to mud in a seaward direction.	C8	Dredged Channel		Unit boundary.
					Approximate unit boundary.

