

7-12-2013

# Tick-Borne Diseases, Maine - 2012

Maine Department of Health and Human Services

Maine Center for Disease Control and Prevention

Follow this and additional works at: [https://digitalmaine.com/mecdc\\_docs](https://digitalmaine.com/mecdc_docs)

---

## Recommended Citation

Maine Department of Health and Human Services and Maine Center for Disease Control and Prevention, "Tick-Borne Diseases, Maine - 2012" (2013). *Center for Disease Control Documents*. 163.  
[https://digitalmaine.com/mecdc\\_docs/163](https://digitalmaine.com/mecdc_docs/163)

This Text is brought to you for free and open access by the Health & Human Services at Digital Maine. It has been accepted for inclusion in Center for Disease Control Documents by an authorized administrator of Digital Maine. For more information, please contact [statedocs@maine.gov](mailto:statedocs@maine.gov).



# Infectious Disease Epidemiology Report

## Tick-Borne Diseases, Maine - 2012



### Background

Tick-borne diseases are a growing concern in the United States. These diseases are transmitted through the bite of an infected tick and can lead to multiple complications.

There are a variety of tick-borne diseases present in Maine. While Lyme disease is the most common tick-borne disease in Maine, there are other diseases of concern to human health including: anaplasmosis, babesiosis, ehrlichiosis, Powassan encephalitis, and Rocky Mountain spotted fever. This surveillance report summarizes the surveillance of tick-borne diseases reported in 2012, excluding Lyme disease. Due to the high incidence of Lyme disease in Maine, it has its own surveillance report which is available on Maine CDC's website at <http://www.maine.gov/dhhs/mecdc/infectious-disease/epi/vector-borne/index.shtml>.

### Anaplasmosis

Anaplasmosis is a bacterial disease transmitted through the bite of an infected deer tick (*Ixodes scapularis*). Signs and symptoms include: fever, headache, malaise, and body aches. Encephalitis or meningitis may occur in rare instances.

### Babesiosis

Babesiosis is a parasitic disease transmitted through the bite of an infected deer tick (*Ixodes scapularis*). Patients may be asymptomatic or may experience symptoms including: extreme fatigue, aches, fever, chills, sweating, dark urine, and anemia. Persons with underlying conditions such as asplenia are at higher risk of severe disease.

### Ehrlichiosis

Ehrlichiosis is a bacterial disease transmitted through the bite of an infected lone star tick (*Amblyomma americanum*). Signs and symptoms include: fever, headache, nausea, rash, and body aches. Encephalitis or meningitis may occur.

### Rocky Mountain Spotted Fever

Rocky Mountain spotted fever (RMSF) is a bacterial disease transmitted through the bite of an infected dog tick (*Dermacentor variabilis*). Signs and symptoms

include: fever, rash, headache, nausea, vomiting, abdominal pain, muscle pain, lack of appetite, and red eyes.

### Methods

All cases of tick-borne diseases in humans are investigated. Standardized case report forms are completed for all cases. Confirmed and probable cases are reported to federal CDC. Cases are classified using CSTE's case definitions for each specific disease.

### Results

A total of 68 confirmed and probable cases of tick-borne disease (excluding Lyme) were reported in 2012. Anaplasmosis was the most commonly reported of these diseases (Table 1).

Table 1: Confirmed & Probable tick-borne diseases - Maine, 2012


Disease	Case Count	Case Rate*
Anaplasmosis	52	3.9
Babesiosis	10	0.8
Ehrlichiosis	3	0.2
RMSF**	3	0.2

\* Case rate per 100,000 persons

\*\*All cases of RMSF are probable

Tick-borne disease cases in Maine have been gradually increasing in number (Figure 1) and expanding in geographic range.

Figure 1: Tick-borne Disease Cases – Maine, 2008 - 2012



## Tick-Borne Diseases – Maine, 2012

Cases of tick-borne diseases are more common in Southern and coastal Maine but are gradually spreading throughout the state (Table 2).

**Table 2: Confirmed and probable tick-borne disease cases, by county – Maine, 2012**

	Anaplasmosis	Babesiosis	Ehrlichiosis	RMSF*
Androscoggin	1			
Aroostook				
Cumberland	10	3		
Franklin				
Hancock	2			1
Kennebec	2	1		
Knox	9	2		1
Lincoln	9		1	
Oxford				
Penobscot	1	1		1
Piscataquis				
Sagadahoc			1	
Somerset	1			
Waldo				
Washington	1			
York	16	3	1	
<b>Maine Total</b>	<b>52</b>	<b>10</b>	<b>3</b>	<b>3</b>

\*All cases of RMSF are probable

### Discussion

Lyme disease is firmly established in Maine, but there are other tick-borne illnesses that are becoming more common. The agents that cause anaplasmosis and babesiosis are transmitted by the same tick that carries Lyme disease, and the numbers of both of these diseases appear to be on the rise. The number of cases of anaplasmosis in the state hit an all-time high of 52 cases in 2012, double the number of cases reported in 2011. There were 4 reported co-infections of Lyme disease and anaplasmosis and 1 reported co-infection of Lyme disease and babesiosis. Fifty percent of the anaplasmosis cases reported a history of tick bite.

Ehrlichiosis is transmitted by a tick that is infrequently found in Maine. Travelers to Southeastern and South-central states may be exposed to this tick much more commonly than they would be in Maine. This disease is steadily moving northward throughout the United States, and is being monitored as an emerging disease in Maine.

RMSF is uncommon in Maine, but the tick that transmits the disease is frequently found here. There are no confirmed cases of RMSF that have been

acquired in Maine; however this is another disease that is considered to be emerging in the state.

### Prevention

To lower the chances of contracting a tick-borne disease, measures should be taken to prevent tick bites both at home and while traveling:

- Using caution in tick-infested areas
- Using EPA-approved repellents on uncovered skin and clothing
- Wearing long sleeved shirts and long pants
- Checking for ticks after being outside
- Removing attached ticks with tweezers or a tick spoon immediately to avoid them becoming engorged
- Using “tick-safe” landscaping such as removing leaf litter, tall grass and brush, creating borders between woods and lawn and discouraging deer with physical barriers

Health care providers are encouraged to consider tick-borne diseases in patients with appropriate clinical presentations. PCR is the preferred method of testing for anaplasmosis, babesiosis, and ehrlichiosis.

All cases of anaplasmosis, babesiosis, ehrlichiosis, Powassan, and Rocky Mountain spotted fever in Maine must be reported within 48 hours by calling 1-800-821-5821, or by faxing reports to 207-287-6865.

Ticks may be submitted for identification free of charge to the Maine Medical Research Institute. Information can be found at:

<http://www.mmcri.org/lyme/submit.html>. Ticks will not be tested for presence of disease.

Additional information about tick-borne diseases can be found at:

- Maine CDC  
<http://www.maine.gov/dhhs/mecdc/infectious-disease/epi/vector-borne/index.shtml>
- Federal CDC  
<http://www.cdc.gov/ticks/index.html>