

**DEPARTMENT OF CONSERVATION
Maine Geological Survey**

Robert G. Marvinney, State Geologist

OPEN-FILE NO. 99-127

Title: *Surficial Geology of the Portsmouth 7.5-minute quadrangle,
York County, Maine*

Author: *Geoffrey W. Smith*

Date: *1999*

Financial Support: Funding for the preparation of this report was provided in part by the U.S. Geological Survey Cooperative Geological Mapping (COGEO-MAP) Program, Cooperative Agreement No. 14-08-0001-A0381.

Associated Maps: Surficial geology of the Portsmouth quadrangle, Open-File 99-96
Surficial materials of the Portsmouth quadrangle, Open-File 98-161

Contents: 5 p. report

Surficial Geology of the Portsmouth 7.5-minute Quadrangle, York County, Maine

Geoffrey W. Smith
Department of Geological Sciences
Ohio University
Athens, Ohio 45701

INTRODUCTION

The Kittery-South Berwick area (Dover East and Portsmouth 7.5-minute quadrangles) is located along and immediately inland from the southwest coast of Maine, approximately 45 miles south of Portland. The area lies within the Seaboard Lowland physiographic province, where altitudes range from sea level to 400 feet above sea level. Most of the area is drained by the Salmon Falls River and its tributaries, which flow to the coast in the vicinity of Kittery and Portsmouth.

The Kittery-South Berwick area and adjacent parts of southwestern Maine are underlain by a variety of igneous intrusive rocks and complexly deformed metamorphic rocks that dip steeply and strike in a general northeast-southwest direction. The dominant structural grain of the bedrock is clearly reflected both in the present topography of the uplands and in the course of many streams that drain the area. This structural control is also evident in the orientation of the many embayments along the present coastline, particularly north of Kittery.

During the last (Wisconsinan) episode of glaciation, ice advanced from the northwest across the area to a terminal position on the continental shelf. Glacial erosion produced a distinct northwest-southeast lineation, superimposed upon the northeast-southwest topographic grain. Streamlined erosional features are common, and several valleys paralleling the direction of ice movement display the effects of erosional deepening and steepening. Glacial deposition resulted in a general reduction of preglacial relief by preferential infilling of valleys. This effect is most pronounced in that portion of the area that experienced late-glacial marine submergence.

Ice retreat, accompanied by marine submergence, progressed rapidly across the area in a general northwesterly direction. End moraines were produced at or near the glacier margin during the period of retreat, and outline the pattern of ice withdrawal.

The first surficial geologic mapping of the Dover East and Portsmouth quadrangles was conducted by Bloom (1960) at a re-

connaisance level (1:62,500; 1:250,000). The area was remapped by J. T. Andrews at scales of 1:24,000 and 1:62,500 in the early stages of the Maine Geological Survey's reconnaissance surficial mapping program. The surficial geology of these quadrangles was revised and mapped in greater detail at a scale of 1:24,000 by G. W. Smith during the 1986 field season (Smith, 1998, 1999).

Information bearing on the stratigraphy and glacial geologic history of this portion of coastal Maine can be found in the following publications: Bloom (1960, 1963) and Smith (1981, 1982, 1984, 1985). Publications by Thompson (1979, 1982), Stuiver and Borns (1975), and Thompson and Borns (1985) provide helpful general references to the glacial geology of the entire coastal zone.

GLACIAL AND POSTGLACIAL DEPOSITS

The general succession of glacial and postglacial deposits in the Kittery-South Berwick area is presented in Figure 1 and Table 1. All glacial deposits in the area are ascribed to the Late Wisconsinan glacial episode. Deposits exposed above the limit of late-glacial marine submergence (which is approximately 200 feet above present sea level) record both glacial advance and retreat, while most sediments exposed below the marine limit are related to deglaciation and the accompanying submergence.

Glacial Till

Glacial till (map units Pt and Ptl) occurs throughout the Kittery-South Berwick area, both above and below the marine limit. The thickness of the till is variable, as is its composition. Below the marine limit, till occurs as a variety of genetic types. Lodgement till, flowtill, and melt-out till comprise the cores of many small moraines. Lodgement till deposited beneath the ice




Figure 1. Generalized glacial stratigraphy for the southern Maine coastal zone (from Smith, 1985.)

sheet forms a blanket deposit over topographic highs, and is inferred to underlie younger deposits in topographic lows. This till is typically a bouldery, gray, compact material, with a silt-sand-clay matrix. Locally, thicker accumulations of lodgement till have been glacially streamlined to form drumlins. Elsewhere, till has been substantially reworked by waves and currents during marine submergence to form **nearshore deposits** (map units Pmn).

TABLE 1. AGE RELATIONSHIPS OF SURFICIAL DEPOSITS IN THE DOVER EAST AND PORTSMOUTH QUADRANGLES*

	Glacial	Glacial Fluvial	Glacial Marine	Fluvial	Wetland	Marine
HOLOCENE			Ha	Hwsm Hws Hwh Hwfm		
			Qst			Pms Pmn
PLEISTOCENE		Pmrs Pp Pm				
LATE WISCONSINAN		Pemc				
	Pt Ptl					

* Units that occur in the Portsmouth quadrangle are shown in boldface type.

End Moraine Complexes

End-moraine complexes (map unit Pemc) have been mapped in several places. In the general area of southwestern Maine, they primarily consist of a variety of end moraines, locally associated with ice-frontal (or marginal) deltas. These deposits can be traced northwestward into the foothills of the White Mountains, where they give way to eskers and kettled valley trains. In their distal portions, the ice-contact deposits commonly intertongue with sand and silt of the marine Presumpscot Formation.

Stratified Deposits

Stratified deposits of sand and gravel that both underlie and intertongue with the Presumpscot Formation are considered to be subaqueous outwash sediments (Smith, 1982, 1985; Thompson, 1982). These materials are not exposed at the ground surface in the study area, and thus are not shown on the geologic map, but they may form an extensive blanket of deposits beneath the younger marine sediments. Many exposures of these materials in nearby areas of southwestern Maine display a variety of distinctly fluvial primary structures. Other deposits have the appearance of subaqueous debris flows. In many cases, the sediments have been severely deformed by thrusting and ice shove, and are complexly interstratified with till and marine silt, sand, and clay.

Glacial-Marine Sediments

Glacial-marine sediments of the **Presumpscot Formation** (map units Pp, Pps, and Pm) occur as a discontinuous cover up to 50 m thick throughout the area of late-glacial marine submergence. The general distribution of the marine sediments was originally mapped by Goldthwait (1949), and the sediments were described in detail and given formational status by Bloom (1960). The marine clay and silt (Pp) is the type Presumpscot Formation described by Bloom. It has a gradational contact with the overlying sandy facies of the marine sediments (unit Pmrs), which is a regressive deposit.

The Presumpscot Formation bears a complex stratigraphic relationship to other glacial sediments below the marine limit. It generally overlies till with sharp contact. In the nearby Berwick-Kennebunk region, subaqueous outwash underlies and intertongues with the Presumpscot Formation, while younger sand and gravel deposits overlie and intertongue with the Presumpscot sediments. Local **shoreline deposits** (map unit Pms) gradationally overlie the Presumpscot Formation and record successive stages of the falling late-glacial sea.

Holocene Deposits

Deposits of Holocene age associated with wetlands, modern streams, and coastal settings have been mapped throughout the study area. In the vicinity of the present coastline, they occur as **salt marshes** (map unit Hwsm). Inland, the wetland deposits include **heaths** (Hwh), **freshwater marshes** (Hwfm), and **swamps** (Hws). Some wetlands are underlain by peat, and are further classified according to the thickness and ash content of the peat (see map explanation). Fluvial **stream-terrace deposits** (map unit Qst) of late-glacial to postglacial age occur locally in stream valleys, at elevations higher than the present-day flood plains. The alluvium on modern flood plains has been mapped as unit Ha.

GLACIAL AND POSTGLACIAL HISTORY

All glacial deposits found in the Kittery-South Berwick area have been ascribed to the most recent (Late Wisconsinan) glacial episode to affect coastal Maine. The orientations of drumlins and other streamlined landforms, as well as of glacial striations, indicate that the last ice to cover the area advanced from the northwest. Divergence from this general trend resulted from large-scale topographic control of the pattern of ice flow.

Withdrawal of Late Wisconsinan ice from its terminal position on the continental shelf was underway between 17,000 and 15,000 years ago, and the ice had retreated across the Gulf of Maine to a position roughly parallel to, but some distance offshore of, the present coastline by 14,000 years ago (Smith, 1985). Dates on shells collected from Great Hill and the Kennebunk landfill site, north of Kittery, suggest that glacial ice may have remained in the vicinity of the coast until 13,200 years ago.

It should be noted, however, that some workers feel that ice had retreated from the coastal area much earlier (Thompson and Borns, 1985).

Two indirect effects of glaciation had a very strong bearing on the character of ice retreat and the deposition of the glacial sediments in this portion of the coastal zone. First, the great weight of the Laurentide continental ice sheet depressed the crust beneath the glacier significantly below its present level throughout the region. Secondly, as glaciers expanded worldwide, water was trapped on land as ice, causing sea level to be lowered by several hundred feet. Later, as the ice began to melt and retreat, water was returned to the ocean and sea level rose immediately. At the same time, the crust began slowly to rebound to its original level. The interaction of these effects resulted in submergence of the entire Maine coastal zone for a period of several hundred years following the retreat of the ice. Furthermore, during its retreat the glacier was grounded on the sea floor, and a complex assemblage of glacial-marine sediments was deposited in the area below the marine limit.

As the glacial ice began to retreat in a general northwesterly direction across the Kittery-South Berwick area, coarse clastic sediments (till, ice-contact stratified drift, subaqueous outwash) accumulated in a narrow zone adjacent to the ice front, while fine sediment (silt and clay of the Presumpscot Formation) was deposited farther away from the ice (Figure 2). Continued ice retreat resulted in the overlap of distal (fine) sediments on proximal (coarse) sediments.

During the period of retreat, at least while the glacier was marine-based, it remained active and continued to advance periodically over short distances. These advances resulted in deformation of previously deposited sediments and construction of minor (DeGeer) moraine ridges. During periods of extended stillstand in the nearby Berwick-Kennebunk area, sediments accumulated at the ice front (or grounding line) to form larger stratified end moraines and deltas.

As ice continued to retreat northwestward across the coastal zone, isostatic rebound began to elevate the land, and relative sea level began to fall. Meltwater streams continued to carry sediment to the ocean as sea level fell. With progressive lowering of sea level, materials deposited during earlier stages of glaciation and deglaciation passed through the wave zone, were eroded, and shed sandy sediment (Pmrs) over the silt and clay (Pp) deposited when sea level was higher. Sandy to silty sediments eroded from till deposits (Pmn) form a veneer over till and bedrock in some parts of the study area. Short-term pauses in the lowering of relative sea level allowed for greater erosion at certain elevations, producing wave-cut scarps and accumulations of beach sediments (Pms). Several of these pauses occurred between the time of maximum marine submergence and offlap to the position of present sea level. Particularly prominent stands of sea level are recorded at elevations of 220-, 200+, 190+, 140+, 100+, 80+, 60+, 40-, and 50+ ft. Inland from the regressing late-glacial shoreline, stream terraces and associated deposits (Qst) began to form along stream valleys.




Figure 2. Glacial marine facies associations for the Maine coastal zone (from Smith, 1984).

Following withdrawal of ice from the coastal zone and the completion of isostatic rebound, eustatic sea level has gradually risen to its present position during Holocene time. Streams are graded to this level and have constructed flood plains. In the coastal zone, marine processes construct sand beaches and develop salt-marsh deposits within the zone of tidal fluctuation. Fine-grained sediments and organic materials (including peat) continue to accumulate in freshwater wetlands.

REFERENCES

Bloom, A. L., 1960, Late Pleistocene changes of sea level in southwestern Maine: Maine Geological Survey, 140 p.
 Bloom, A. L., 1963, Late Pleistocene fluctuations of sea level and postglacial crustal rebound in coastal Maine: American Journal of Science, v. 261, p. 862-879.
 Goldthwait, L., 1949, Clay Survey - 1948: Maine Development Commission, Report of the State Geologist, 1947-1948, p. 63-69.
 Prescott, G. C., and Drake, J. A., 1962, Records of selected wells, test holes, and springs in southwestern Maine: U.S. Geological Survey, Maine Basic Data Report No. 1, Ground water series: southwestern area, 35 p.
 Smith, G. W., 1981, The Kennebunk glacial advance: A reappraisal: Geology, v. 9, p. 250-253.
 Smith, G. W., 1982, End moraines and the pattern of last ice retreat from central and south coastal Maine, in Larson, G. J., and Stone, B. D. (eds.), Late Wisconsinan glaciation of New England: Kendall/Hunt Publishing Co., Dubuque, Iowa, p. 195-209.

Smith, G. W., 1984, Glaciomarine sediments and facies associations, southern York County, Maine, in Hanson, L. S. (ed.), Geology of the coastal lowlands, Boston, MA to Kennebunk, ME: New England Intercollegiate Geological Conference, p. 352-369.
 Smith, G. W., 1985, Chronology of Late Wisconsinan deglaciation of coastal Maine, in Borns, H. W., Jr., LaSalle, P., and Thompson, W. B. (eds.), Late Pleistocene history of northeastern New England and adjacent Quebec: Geological Society of America, Special Paper 197, p. 29-44.
 Smith, G. W., 1998, Surficial materials of the Portsmouth quadrangle, Maine: Maine Geological Survey, Open-File Map 98-162.
 Smith, G. W., 1999, Surficial geology of the Portsmouth quadrangle, Maine: Maine Geological Survey, Open-File Map 99-96.
 Stuiver, M., and Borns, H. W., Jr., 1975, Later Quaternary marine invasion in Maine: its chronology and associated crustal movement: Geological Society of America, Bulletin, v. 86, p. 99-104.
 Thompson, W. B., 1979, Surficial geology handbook for coastal Maine: Maine Geological Survey, 68 p.
 Thompson, W. B., 1982, Recession of the Late Wisconsinan ice sheet in coastal Maine, in Larson, G. J., and Stone, B. D. (eds.), Late Wisconsinan glaciation of New England: Kendall/Hunt Publishing Co., Dubuque, Iowa, p. 211-228.
 Thompson, W. B., and Borns, H. W., Jr., 1985, Till stratigraphy and late Wisconsinan deglaciation of southern Maine: A review: Geographie physique et Quaternaire, v. 39, no. 2, p. 199-214.
 Tolman, A. L., Tepper, D. H., Prescott, G. C., and Gammon, S. O., 1983, Hydrogeology of significant sand and gravel aquifers: northern York and southern Cumberland Counties, Maine: Maine Geological Survey, Open-File Report 83-1.

APPENDIX A. TABULATION OF DATA FROM PRESCOTT AND DRAKE (1962)

Records of Selected Wells

Well No.	Location	(altitude in ft)	Depth of Well (ft)	Aquifer Material
218	Kittery	(40)	96	Bedrock
433	Kittery	(40)	69	Bedrock
435	Kittery	(40)	59	Bedrock
440	Eliot	(50)	90	Bedrock
441	Eliot	(50)	95	Bedrock
444	Kittery	(40)	52	Bedrock
621	Eliot	(80)	26	
622	Eliot	(50)	17	
624	Eliot	(40)	45	

Records of Selected Springs

Well No.	Location	(altitude in ft)	Depth of Well (ft)	Aquifer Material
139	Eliot	(70)		Sand/gravel

**APPENDIX B. TABULATION OF DATA FROM
MAINE DEPARTMENT OF TRANSPORTATION SOILS REPORT 85-101**

Boring CB-59-84	5 ft	Brown silty sand and gravel; possible ledge
Boring CB-48-84	5 ft	Brown sand and gravel; possible ledge
Boring CB-44-84	5 ft	Brown sand and gravel; apparent ledge
Boring CB-60-84	5 ft	Brown sand and gravel; 100% ledge
Boring CB-63-84	2 ft	Brown sand and gravel; ledge
Boring CB-43-84	5 ft	Brown sand and gravel
Boring CB-64-84	3 ft	Brown sand and gravel; ledge
Boring CB-61-84	3 ft	Brown sand and gravel; ledge
Boring CB-62-84	3 ft	Brown sand and gravel; ledge
Boring CB-65-84	6 ft	Brown sand and gravel; ledge (?)
Boring CB-47-84	1 ft	Brown sand and gravel; ledge (?)
Boring CB-45-84	5 ft	Brown sand and gravel; ledge
Boring CB-46-84	8+ ft	Brown sand and gravel