


Bedrock Geology



Newbury Neck Quadrangle, Maine

Bedrock geologic mapping by


Douglas N. Reusch
John P. Hogan

Digital cartography by:
Susan S. Tolman

Robert G. Marvinney
State Geologist

Cartographic design and editing by:
Robert D. Tucker
Henry N. Berry IV

Funding for the preparation of this map was provided in part by the U.S. Geological Survey STATEMAP Program, Cooperative Agreement No. 00HQAG007.



Maine Geological Survey

Address: 22 State House Station, Augusta, Maine 04333
Telephone: 207-287-2801 **E-mail:** mgs@state.me.us
Home page: <http://www.state.me.us/doc/nrmc/nrmc.htm>

Open-File No. 02-162

2002

For additional information,
see Open-File Report 03-92.

GEOLOGIC HISTORY

ORIGIN OF THE STRATIFIED ROCKS

The Ellsworth Schist, which underlies most of the map area, consists of a pile of weakly metamorphosed sedimentary and igneous rocks 4 to 7 kilometers thick. The majority of the Ellsworth Schist is a metamorphic rock called schist, which gives its name to the formation. Schist originated as sediments that were eroded from nearby volcanic materials and deposited in an ocean in Cambrian time (see Geologic Time Scale below). The volcanic materials probably included lava flows and deposits of rock fragments of various sizes. Most of the schists are very thinly layered rocks and represent fine-grained sediments such as mud and silt that have since been changed by heat and pressure into schist. A few quartzite beds (Photo 1) were originally layers of sand. A thin conglomeratic unit at the head of Morgan Bay (Cems) was originally a layer of pebbly gravel. These layers indicate the sediments originally accumulated in a basin near land. A part of the Ellsworth Schist that has more quartzite layers than other parts is indicated on the map by the symbol Cem.

Thin sheets of igneous rock are widespread in the Ellsworth Schist. Igneous rocks originate from molten rock, or magma, that cools and solidifies. Most of the igneous rocks in the Ellsworth Schist are of two types, a dark greenish gray rock called greenstone (Photo 2) and a

white to cream-weathering rock called rhyolite (Photo 3). Greenstone forms from dark gray to black lava flows or volcanic ash (of basaltic composition) which has been altered through chemical reactions to produce a rock rich in the dark green mineral chlorite. Rhyolite is of a chemical composition different from greenstone, the same composition as granite. But in contrast to slowly-cooled granite, rhyolite cools rapidly from the molten state to form a uniform, hard, crystalline rock with very small mineral grains. The association of greenstone and rhyolite, together with other chemical characteristics determined by laboratory analysis, suggests the igneous rocks of the Ellsworth Schist formed in a continental rift (see Stewart, 1998). Most of these igneous rocks formed when lava erupted at the earth's surface, in some cases violently, during the time the sediment was accumulating. A few igneous sheets, however, solidified more slowly below the ground surface as indicated by their larger grain size and crosscutting relationships (Photo 4). The unit Cer at the eastern side of the map is composed largely of rhyolite produced during a larger volcanic episode. The unit Cee at the northeast corner of the map, though poorly exposed, appears to contain a higher proportion of greenstone than does the Ellsworth Schist elsewhere in the map area.



Photo 1. Thin layer of white quartzite in gray schist. Such well-defined quartzite layers are rare in most of the Ellsworth Schist. (East shore of Newbury Neck, 4000 feet north of Curtis Cove.)



Photo 2. Greenstone sheet four inches thick, parallel to foliation in the schist. View looking north. (Weymouth Point, mouth of Union River.)



Photo 3. White rhyolite. The dark fragment within the rhyolite is a fragment of a different rock type. This demonstrates that the rhyolite layer is an accumulation of rock fragments, formed by a violent volcanic eruption. (East shore of Newbury Neck, 6000 feet north of Curtis Cove.)



Photo 4. Angular chunk of rhyolite in greenstone. The light and dark spots in the greenstone are relatively large mineral grains that indicate this igneous sheet solidified underground. (North end of Morgan Bay.)

ROCK DEFORMATION AND RELATED METAMORPHISM

The Ellsworth Schist has been deformed at least three times, producing a highly contorted rock structure. In the main stage of deformation, wet sediments together with the interlayered volcanic materials were squeezed from the sides, piling up toward the northwest. Distorted thin layers show that the rocks were crumpled into folds inclined toward the northwest (Photo 5). Curved shapes, called shear bands, also indicate the upper part of the formation was pushed toward the northwest over the lower part (Photo 6). This deformation affected the entire mass of the Ellsworth Schist. The most pervasive feature of every outcrop is the main foliation, a microscopic alignment of flat mineral grains giving the rock an overall sheet-like structure. On the face of foliation surfaces, mineral grains have been stretched into lines to produce a mineral lineation. This lineation may be defined by elongate quartz grains or trails of broken minerals such as feldspar and pyrite. In most places the lineation is oriented in a northwest-southeast direction as represented by symbols on the map.

The minerals that comprise the main foliation, chlorite and white mica, are metamorphic minerals that form at elevated temperatures, typically 350 to 400 degrees Centigrade. It is due to this heat that the rocks could be deformed by folding rather than by breaking. Also during the heating, milky white quartz veins formed in the schist. These veins are scattered throughout the formation and were deformed along with the rest of the rock (Photo 7). The age of this deformation and metamorphism is not known very precisely; it must be younger than the Cambrian age of the Ellsworth Schist which is not deformed, and older than the Devonian granite of Blue Hill which is not deformed.

The main foliation and lineation of the Ellsworth Schist were affected by a variety of later folds (Photo 8). Some of them appear to indicate movement towards the southeast, in a direction opposite to the earlier main stage movement toward the northwest. Late folds are particularly well-displayed at High Head in a zone of complex deformation 500 meters wide.

Following the two stages of folding, a few late brittle faults occurred in the area. The rocks at Weymouth Point are broken along

steeply inclined faults approximately parallel to the length of Patten Bay (Photo 9). The orientation of these minor faults suggests that an old fault trace may lie beneath Patten Bay. Another piece of circumstantial evidence is the presence of metamorphic biotite (black mica) south of Patten Bay and its absence north of the bay. While this difference might be just the original metamorphic pattern, it might also be explained by a fault, as shown on the map by the truncated metamorphic isograd line. Clearly, this large fault is not proven to exist; it is inferred from the minor faults observed in outcrop. The faults may be related to an extension of the crust in a northwest-southeast direction. The age of the faults is probably younger than the Devonian granite of Blue Hill and may be Mesozoic.



Photo 5. Lower part of photo shows quartzite layers and an early foliation deformed into folds. The asymmetry of the folds indicates the top moved towards the northwest (left). Upper part of photo shows the main foliation is also inclined to the left. (North shore of Western Bay, opposite Alley Island. View toward the northeast.)



Photo 6. White quartz veins in schist deformed by horizontal shear bands. The sense of curvature of the quartz vein tails indicates top-to-the-northwest (left) movement. The quartz vein to the right of the penny is very tightly folded. (West shore of Union River Bay south of Patten Bay. View toward the northeast.)



Photo 7. Typical shoreline exposure of contorted Ellsworth Schist, looking south at High Head toward Goose Rock. White quartz veins in foreground formed during metamorphism. (South end of Newbury Neck.)




Photo 8. Small, late folds deform layering and foliation in Ellsworth Schist. These particular ones indicate movement to the north, though most late folds are asymmetric toward the south. (Goose Rock, south end of Newbury Neck.)




Photo 9. Sheet of greenstone parallel to the main foliation in schist. A late, brittle fault to the right of the notebook has offset the left (southeasterly) block downward by more than a foot with respect to the equivalent rocks on the right. (West shore of Union River, north of Weymouth Point.)




Photo 10. Large mineral grains in pegmatite. White = quartz, off-white = feldspar, black = tourmaline. (Webber Cove, west side of Morgan Bay.)




Photo 11. Andalusite crystals in Ellsworth Schist. This mineral grew in response to heating of rocks outside the Blue Hill granite. (North end of Morgan Bay.)

All the metamorphic and intrusive bedrock in the map area originally formed at depth in the earth. In order for these rocks to be exposed at the surface now, a significant amount of overlying rock must have been eroded in the hundreds of millions of years since the Devonian Period. The latest increment of erosion occurred during the last Ice Age.

when a continental glacier extended across Maine and ground down some of the bedrock surface. When the last glacier melted from the Maine coast, about 14,000 years ago, it deposited a layer of sediment burying most of the bedrock in the area. These sediments are described on surficial geologic maps, available from the Maine Geological Survey.

INTRUSIVE ROCKS

Mesozoic or Paleozoic

Dikes of dark gray to light gray intrusive rock. Mainly diabase but also includes rocks of intermediate composition. Letter symbols indicate composition: m=mafic, i=intermediate. Line symbols are drawn parallel to strike and annotated with dip if known. (Dip unknown, Inclined, Vertical)

Devonian

Granite of Blue Hill. Gray, medium-grained equigranular muscovite-bearing biotite granite. Rb-Sr whole-rock age is 374 ± 10 Ma (Brooks, 1976, recalculated using ^{87}Rb decay constant of $1.42 \times 10^{-11}/\text{yr}$).

Dge Coarse-grained equigranular biotite \pm muscovite syenogranite.

Dgs Coarse-grained seriate biotite \pm muscovite syenogranite.

Dgl Fine-grained to aplitic tourmaline-muscovite leucogranite. In Webber Cove, at the top of sill is quartz-feldspar-tourmaline-muscovite pegmatite. Scattered clots of tourmaline.

Devonian-Silurian

Cadillac Mountain Intrusive Complex

DScg Cadillac Mountain Granite. Pink to greenish-gray, coarse-grained hornblende \pm biotite granite. U-Pb zircon age of 419 ± 2 Ma reported by Seaman and others (1995).

DSgd Gabbro-diorite.

Late Cambrian (?)

Little-deformed massive vesicular greenstone. Possible feeders for Cassin Volcanics. Occurs as dikes at Weymouth Point and High Head. Dip indicated if known.

STRATIFIED ROCKS

Middle Cambrian

Ce **Ellsworth Schist.** Quartz-muscovite-chlorite schist interstratified with lesser metamorphosed bimodal igneous rocks. Schists are typically dark green and light-weathering and consist of anastomosing, variably thick (1-20 mm), fine-grained, equigranular quartz-rich laminations and muscovite-chlorite lenses and films. Most outcrops are generally massive, without discernible bedding, and compositions range from phyllosilicate-rich through quartz-rich. In contact aureole of Blue Hill pluton, pelitic beds contain fine-grained biotite, cm-sized andalusite, and cordierite(?). Formation contains rare medium-bedded quartz-feldspar sandstones and impure quartzites. Ubiquitous sheets of metamorphosed igneous rocks 10 to 200 cm in thickness comprise greenstones (meta-basalts) and meta-rhyolites. Abundant greenstone sheets are typically fine-grained, massive, and contain sparse feldspars and patches of epidote. Most greenstones are considered to be extrusive, but coarse-grained foliated gabbro with chilled margins occurs in sills between the head of Morgan Bay and Browns Point. Rhyolites are light gray, weather cream to white, and typically contain disseminated pyrite cubes a few mm across. Most rhyolites are inferred to be pyroclastic deposits, including white-weathering crystal tuffs and massive white vitric tuffs with sparse 10 cm-sized flattened pumice fragments.

Cee **Egypt Member.** Dark green feldspar porphyroblastic schist. Fine-grained biotite reported in thin sections (McGregor, 1964). Extensively exposed in large boulders on east shore of Union River.

Cem **Morgan Bay Member.** Chlorite schist with sparse thin to medium beds of quartzite.

Cems **Smelt Brook Conglomerate.** Conglomerate bed 30 cm thick near head of Morgan Bay. Contains scattered cm-sized partially flattened pebbles of felsic igneous rocks.

Cer **Rhyolite of Goose Cove.**

EXPLANATION OF SYMBOLS

- Stratigraphic or intrusive contact. (Well-defined, Approximately located, Inferred) Dip indicated if known.
- Inferred fault contact.
- Axial trace of antiform.
- Axial trace of synform.
- Contact metamorphic isograd. Hatchures toward higher grade with index mineral indicated. (Biotite, Andalusite)
- Outcrop
- Outcrop of massive (non-foliated) granite.
- Strike and dip of igneous layering (in Cadillac Mountain Intrusive Complex).
- Strike and dip of felsic dike (in Devonian granite).
- Strike and dip of bedding. (Horizontal, Inclined)
- Strike and dip of igneous sheet. Most are inferred to be volcanic or volcanoclastic layers, except sills at head of Morgan Bay and Browns Point. (Horizontal, Inclined)
- Strike and dip of main foliation, defined by preferred orientation of phyllosilicates. (Horizontal, Inclined)
- Trend and plunge of mineral lineation, defined by elongate quartz, train of broken feldspar or pyrite. (Horizontal, Plunging)
- Trend and plunge of crenulation intersection lineation on foliation surface. (Horizontal, Plunging)
- Main generation folds. Generally asymmetric, tight to isoclinal ductile folds typically a few centimeters in wavelength.
- Hinge line, asymmetry unknown. (Horizontal, Plunging)
- Hinge line, sense of asymmetry indicated. (Clockwise, Counterclockwise, Neutral)
- Late generation folds. Symmetric to asymmetric folds deform foliation and lineation. Generally more open than main generation folds. (Horizontal, Plunging, Clockwise asymmetry, Neutral asymmetry)
- Strike and dip of joint. (Vertical, Inclined)
- Location of photo shown in sidebar.

REFERENCES

- Brooks, D. G., 1976. Geochronologic contributions to stratigraphic interpretation and correlation in the Penobscot Bay area, eastern Maine. In: Page, L. R. (editor), Contributions to the stratigraphy of New England: Geological Society of America, Memoir 148, p. 129-145.
- Gilman, R. A., and Chapman, C. A., 1988. Bedrock geology of Mount Desert Island, in Gilman, R. A., Chapman, C. A., Lovell, T. V., and Borns, H. W., Jr., The geology of Mount Desert Island: a visitor's guide to the geology of Acadia National Park: Maine Geological Survey, Bulletin 38, map scale 1:50,000.
- McGregor, J. D., 1964. Geology of the Ellsworth quadrangle and vicinity, Maine: Ph.D. dissert., University of Illinois, Urbana, 116 p.
- Seaman, S. J., Wobus, R. A., Wiebe, R. A., Lubick, N., and Bowring, S. A., 1995. Volcanic expression of bimodal magmatism: The Cradbury Island-Cadillac Mountain Complex, coastal Maine. Journal of Geology, v. 103, p. 301-311.
- Stewart, D. B., 1998. Geology of northern Penobscot Bay, Maine: U.S. Geological Survey, Miscellaneous Investigation Series Map I-2551, scale 1:62,500.
- Precambrian time
- * In millions of years before present.

GEOLOGIC TIME SCALE

Geologic Age	Absolute Age*
Cenozoic Era	0-66
Mesozoic Era	Cretaceous Period 66-144 Jurassic Period 144-208 Triassic Period 208-245
Paleozoic Era	Permian Period 245-286 Carboniferous Period 286-360 Devonian Period 360-418 Silurian Period 418-443 Ordovician Period 443-495 Cambrian Period 495-545
Precambrian time	Older than 545