


SURFICIAL GEOLOGY
OF THE
EDMUNDSTON
1° X 2° QUADRANGLE, MAINE



Compiled By :
Thomas V. Lowell, Maine Geological Survey

Area Compilers and Contributors :
Andrew N. Genes, University of Massachusetts, Boston
Thomas V. Lowell, Maine Geological Survey

Maine Geological Survey
DEPARTMENT OF CONSERVATION
Augusta, Maine 04333
Walter A. Anderson, State Geologist
1987
OPEN-FILE NO. 87-17



COMPILATION RESPONSIBILITY



Funding for the preparation and publication of this map was provided by a grant from the U.S. Department of Energy (grant no. DE-FG02-81NE46640) and the Maine Geological Survey.

EXPLANATION

Geologic Unit	Materials	Topography	Origin
a Stream alluvium (includes Holocene flood plain, stream terrace, and alluvial fan deposits)	Sand, gravel, and silt.	Flat to gently sloping on flood plains and stream terraces; gently to moderately sloping on alluvial fans.	Deposited on flood plains and stream beds by postglacial streams.
s Swamp, marsh, and bog deposits (includes both fresh-water and salt-water marshes)	Peat, muck, clay, silt, and sand.	Flat.	Formed by accumulation of sediments and organic material in depressions and other poorly drained areas.
b Beach deposits	Sand and gravel.	Gently to moderately sloping, with low ridges and mounds.	Includes beach sediments formed by wave and current action, and sand dunes derived from these deposits.
eb Emerged beach deposits	Sand and gravel.	Low ridges or sloping surfaces. May be associated with wave-cut benches on hillsides.	Formed by wave erosion of till or other materials during the late-glacial marine submergence of parts of southern Maine.
e Eolian deposits	Sand.	Dune ridges and mounds, or blanket deposit that conforms to surface of underlying unit.	Windblown sand. Derived from wind erosion of glacial sediments and deposited in late-glacial to postglacial time.
L Lake-bottom deposits	Silt, clay, and sand. Commonly well stratified, and may be rhythmically bedded.	Flat to gently sloping except where dissected by modern streams.	Composed of sediments that washed out of late Wisconsinan glacial ice and accumulated on the floors of glacial lakes. Map unit may also include a few non-glacial lake deposits.
m Glaciomarine deposits (fine-grained facies)	Silt, clay, sand, and minor amounts of gravel. Commonly a clayey silt (the Presumpscot Formation). Sand is dominant in some places, but may be underlain by finer grained sediments. Locally fossiliferous. Map unit includes small areas of till and other units that are not completely covered by marine sediments.	Flat to gently sloping except where dissected by modern streams. Commonly has a branching network of steep-walled stream gullies.	Composed of glacial sediments that accumulated on the ocean floor. Formed during the late-glacial marine submergence of lowland areas in southern Maine.
ms Glaciomarine deposits (coarse-grained facies)	Sand, gravel, and minor amounts of silt.	Flat to moderately sloping. Steeper on ice-contact slopes and delta fronts. May be kettled where deposited over stagnant ice blocks.	Deposited where glacial meltwater streams and currents entered the sea. Includes glaciomarine deltas, subaqueous kames and fans (subaqueous outwash), and outwash that prograded into shallow marine waters and locally covered earlier glaciomarine silt and clay deposits.
go Glacial outwash deposits	Sand and gravel.	Flat to gently sloping. Steeper on ice-contact slopes and delta fronts. May be kettled where deposited over stagnant ice blocks.	Deposited by meltwater streams in front of the receding late Wisconsinan ice margin. Includes non-marine outwash plains, deltas, and fans.
g Ice-contact glaciofluvial deposits (exclusive of eskers)	Sand, gravel, and silt.	Flat-topped kame terraces and deltas which are locally kettled and bounded by steep sides, or hummocky terrain with numerous kames and kettles.	Deposited by meltwater streams adjacent to stagnant glacial ice.

Geologic Unit	Materials	Topography	Origin
ge Eskers	Gravel and sand. May include minor amounts of till. Portions of many eskers below the marine limit are partly or entirely buried by glaciomarine deposits.	Individual or multiple ridges. Complex eskers may have anastomosing patterns and be gradational with other types of ice-contact deposits.	Chiefly deposited by meltwater streams flowing in tunnels within or beneath the late Wisconsinan ice sheet. Map unit also includes small undifferentiated areas of units "g" and "go".
sm Stagnation moraine	Mostly till, but also includes variable percentages of undifferentiated sand and gravel.	Undulating topography with local hummocks and ridges.	Deposited during the dissipation of stagnant glacial ice.
em End moraines	Till or sand and gravel. May be very bouldery. Commonly interbedded with or overlain by glaciomarine sediments in areas that experienced late-glacial marine submergence. Only the largest end moraines and some dense clusters of smaller ones are shown here as a separate unit (em). Elsewhere, short lines mark the crests of moraine ridges, which are locally so numerous that only selected individuals are represented.	Ridges. Commonly arcuate, discontinuous, and in groups. May be multi-crested and hummocky. Size ranges 1-10 m high, 5-200 m wide, and 30 m to over 10 km long.	Deposited in the marginal zone of the late Wisconsinan ice sheet, by glacial ice and/or meltwater flowing out of the ice.
rm Ribbed moraine	Till is the principal constituent, but stratified sediments are present in some of the deposits.	Numerous hummocks and short sub-parallel ridges which typically occur in lake basins and other lowland areas.	Origin uncertain. Deposited either at the margin of or beneath the late Wisconsinan ice sheet.
t Till	Heterogeneous mixture of sand, silt, clay and stones. May include many boulders. Generally massive, but in many places contain beds and lenses of variably washed and stratified sediments.	Generally a blanket deposit that conforms to the underlying bedrock topography. Also forms drumlins and other glacially streamlined hills.	Deposited directly by glacial ice.
Thin drift	Area of many bedrock outcrops and/or thin surficial deposits (generally less than 3 m thick). The type of surficial material is known or inferred.	Topography of these areas reflects the configuration of the bedrock surface and ranges from smooth undulating hills to knobby terrain and high mountains.	Commonly the result of non-deposition of glacial sediments, but the surficial materials in some coastal areas have been largely removed by marine erosion in late-glacial time.
ldu Thin drift, undifferentiated	Area of many bedrock outcrops and/or near-surface bedrock where the surficial materials have not been mapped.	Same as other thin-drift areas.	Same as other thin-drift areas.
rk Bedrock	Area of extensive bedrock outcrop, or where the bedrock has only a thin cover of soil and vegetation. Surficial deposits are essentially absent. Particularly common on the ridge crests and steeper slopes of mountainous areas.	Hilly to mountainous terrain.	Same as the thin-drift areas.

GEOLOGIC SYMBOLS

	Contact	Boundary between adjacent map units.
	Moraine ridge	Lines mark the crests of individual and moraines. Symbol also is used in conjunction with unit rm to show orientation of drift ridges of uncertain origin.
	Glacial striation locality	Includes striations, grooves, crag-and-tails, and other types of ice-flow indicators on bedrock outcrops. Dot indicates point of observation. Arrow-head is omitted where ice-flow direction is uncertain. Flange indicates older trends.
	Glacially streamlined landform	Symbol shows long-axis orientation of drumlins, fluted till ridges, roches moutonnees, and other hills that have been elongated parallel to the flow of glacial ice.
	Cirque	Steep-walled, semicircular bedrock basin formed by glacial erosion in high mountainous areas.
	Meltwater channel	Channel eroded by glacial meltwater stream. Arrow indicates known or inferred direction of stream flow.
	Glaciomarine delta	Number indicates surveyed altitude (in feet) of contact between topset and foreset beds, or of meltwater channel on delta surface, which approximately marks position of sea level in late-glacial time.
	Glaciolacustrine delta	Number indicates approximate altitude (in feet) of former glacial-lake surface.
	Delta of uncertain origin	Delta formed near limit of late-glacial marine submergence. Number indicates approximate altitude (in feet) of contact between topset and foreset beds.

SITES OF SPECIAL INTEREST

- Location of special site

This list includes locations of important stratigraphic sections of Pleistocene deposits in Maine, and places where good examples of certain glacial features can be seen. The sites were selected partly on the basis of accessibility, ease of observation, and relative permanence. Some features, such as eskers and Doherty moraines, are so numerous that only a few of the best examples are included here.

Site	Town	Name/Description	Principal References
1	St. Francis	Golden Rapids till section; multiple-till exposure with interbedded waterlain sediments.	Becker (1); Mawson, Genes, and Brewer (17)
2	Van Buren	Gravel pit at Parent; exposure showing till overlying glaciofluvial gravel (see of several such localities in this part of the St. John River valley).	Kite (9)

RADIOCARBON-DATED SITES

Explanation of symbols used to designate sites on the map:

- Material in place between late Wisconsinan tills.
- Material that predates or is contemporaneous with the advance of the late Wisconsinan ice sheet.
- Material that postdates or is contemporaneous with the recession of the late Wisconsinan ice sheet.
- Material that approximately dates the onlap of the sea during the late-glacial transgression.
- Material that approximately dates the offlap of the sea.

Site	Name/Town	Date (yr B.P.)	Laboratory No.	Material	Reference
A	Rideout Pond (T19 R12 WELS)	10,385 ± 140	SI-4658	organic material in lake-bottom sediments	Davis and Jacobson (2)
B	Fort Kent	>38,000	W-2372	wood from drilled well	Rubin (18)
C	Green River, New Brunswick	10,220 ± 350	I(GSC)-2	basal peat, underlain by lake-bottom sediments	Kite (9)

SOURCES OF GEOLOGIC INFORMATION

- Becker, D. A., 1962, Late Wisconsinan stratigraphy, upper St. John River, northwestern Maine: M.S. thesis, SUNY, Buffalo, 95 p.
- Davis, R. B., and Jacobson, G. W., Jr., 1963, Late-glacial and early Holocene landscapes in northern New England and adjacent areas of Canada: Quart. Res., v. 22.
- Genes, A. N., 1978, Reconnaissance surficial geology of the Square Lake (15°) quadrangle, Maine: Maine Geol. Surv., Open-File Map 78-5.
- Genes, A. N., 1978, Reconnaissance surficial geology of the Stookholm (15°) quadrangle, Maine: Maine Geol. Surv., Open-File Map 78-6.
- Genes, A. N., 1981, Reconnaissance surficial geology of the Eagle Lake (15°) quadrangle, Maine: Maine Geol. Surv., Open-File Map 81-12.
- Genes, A. N., 1981, Reconnaissance surficial geology of the Fort Kent (15°) quadrangle, Maine: Maine Geol. Surv., Open-File Map 81-14.
- Genes, A. N., 1981, Reconnaissance surficial geology of the St. Francis (15°) quadrangle, Maine: Maine Geol. Surv., Open-File Map 81-15.
- Genes, A. N., unpublished reconnaissance data.
- Kite, J. S., 1979, Postglacial geologic history of the middle St. John River valley: M.S. thesis, Univ. of Maine, Orono, 136 p.
- Kite, J. S., unpublished reconnaissance data.
- Lowell, T. V., 1986, Reconnaissance surficial geology of the Allagash (15°) quadrangle, Maine: Maine Geol. Surv., Open-File Map 86-41.
- Lowell, T. V., 1986, Reconnaissance surficial geology of the Beau Lake (15°) quadrangle, Maine: Maine Geol. Surv., Open-File Map 86-42.
- Lowell, T. V., 1986, Reconnaissance surficial geology of the Little East Lake (15°) quadrangle, Maine: Maine Geol. Surv., Open-File Map 86-44.
- Lowell, T. V., 1986, Reconnaissance surficial geology of the Rocky Brook (15°) quadrangle, Maine: Maine Geol. Surv., Open-File Map 86-45.
- Lowell, T. V., 1986, Reconnaissance surficial geology of the Rocky Mtn. (15°) quadrangle, Maine: Maine Geol. Surv., Open-File Map 86-46.
- Lowell, T. V., unpublished reconnaissance data.
- Mawson, A. A., Genes, A. N., and Brewer, T., 1980, Pleistocene geology of northeastern Maine, in Burns, H. W., Jr., LaSalle, P., and Thompson, W. B., (eds.), Late Pleistocene History of Northeastern New England and Adjacent Quebec, Geol. Soc. Amer., Special Paper 197, p. 59-70.
- Rubin, M., 1977, U.S. Geol. Surv., Radiocarbon Laboratory Report No. WB-2372.

INDEX TO
SOURCES OF GEOLOGIC INFORMATION

