



Surficial Geology



SOURCES OF INFORMATION

Surficial geologic mapping of the Purgatory quadrangle was conducted by Carol T. Hildreth during the 2002 field season. Funding for this work was provided by the U. S. Geological Survey STATEMAP program and the Maine Geological Survey, Department of Conservation.



NOTE: A thin, discontinuous layer of windblown sand and silt, generally mixed with underlying glacial deposits by first action and bioturbation, is present near the ground surface over much of the map area but is not shown.

Marine fan deposits (Pleistocene) - Composed mostly of stratified sand and gravel deposited as submarine fans at the glacier margin. Locally underlain by unit Pp. Includes sediments reworked by marine nearshore processes during regression of the sea.

Artificial fill - Man-made. Material may vary from natural sand and gravel to quarry waste or sanitary landfill, including highway and railroad embankments. This material is mapped only where it can be identified using the topographic contour lines or where actually observed. Minor artificial fill is present in virtually all developed areas of the quadrangle. Thickness of fill varies.

Till (Pleistocene) - Light- to dark-gray, nonsorted to poorly sorted mixture of clay, silt, sand, pebbles, cobbles, and boulders; a predominantly sandy diamicton containing some gravel. Generally older than other deposits and may underlie them. Thickness varies and generally is less than 6 m (20 ft), but may be more than 30 m (100 ft) in areas of streamlined topography. Many streamlined hills in this area are bedrock-cored.

Stream alluvium (Holocene) - Sand, silt, gravel, and muck in flood plains along present rivers and streams. As much as 3 m (10 ft) thick. Extent of alluvium indicates most areas flooded in the past that may be subject to future flooding. In places, this unit is indistinguishable from grades into, or is interbedded with wetland deposits (Hw).

Thin glacial drift (Pleistocene) - Areas with generally less than 3 m (10 ft) of drift covering bedrock. Till overlies bedrock on hillslopes and ridge crests; Presumpscot Formation silt and clay is present in depressions; and nearshore deposits overlie till. Presumpscot Formation and bedrock on hillslopes and at the base of these slopes.

Wetland deposit (Holocene) - Muck, peat, silt, and sand deposited in poorly drained areas. Generally 0.5 to 3 m (1 to 10 ft) thick, but may be thicker in bogs. In places, this unit is indistinguishable from grades into, or is interbedded with wetland deposits (Hw).

Bedrock exposures - Not all observed individual outcrops are shown on the map. Gray dots indicate observed outcrops; ruled pattern indicates areas of abundant exposures and areas where surficial deposits are generally less than 3 m (10 ft) thick. Mapped in part from aerial photography, soil surveys (McEwen, 1970; Faust and LaFlamme, 1978), and previous geologic (Thompson and Smith, 1977; Thompson and Borns, 1985) and materials maps (Locke, 1999).

Eolian deposit (Late Pleistocene) - Fine- to medium-grained, well-sorted sand. Found as small dunes on a variety of older glacial deposits. Deposited after late-glacial sea level regression from the area and left fine-grained marine sediments exposed to wind erosion and transport before vegetation established itself and anchored the deposits. Thickness varies from 0.5 to 2 m (1 to 6 ft).

Glaciomarine delta topset-foreset contact elevation (in feet above modern-day sea level; from Thompson and others, 1989).

Marine nearshore deposits (Pleistocene) - Sand, silt, mud, and minor gravel. Consists of reworked till and glaciomarine sediments, redistributed by marine currents and wave action as sea level fell during late-glacial time. May include shoreline, beach, and dune deposits in places. As much as 3 m (10 ft) thick.

Presumpscot Formation: Glaciomarine bottom deposits (Pleistocene) - Silt and clay with local sandy beds. Consists of late-glacial sea-floor deposits. Commonly lies beneath surface deposits of unit Pmn. In places, may be overlain by unmaped thin dune deposits. As much as 21 m (70 ft) thick.

Marine delta deposits (Pleistocene) - Mostly sorted and stratified sand and gravel. Consists of glaciomarine delta deposits, part of which may be ice-contact deposits. Deposits are graded to surface of late-glacial sea and are distinguished by flat top and foreset and topset beds. Deltas have been assigned unique geographic names listed below. Thickness varies from 0 to 30 m (0-100 ft).

Glaciomarine delta topset-foreset contact elevation (in feet above modern-day sea level; from Thompson and others, 1989).

Pmdg -- Grant Cemetery delta

Pmdg -- Grant Cemetery delta

Pmdg -- Grant Cemetery delta

Pmdg -- Grant Cemetery delta

----- **Contact** - Boundary between map units, approximately located.

--- **Channel** - Eroded by glacial meltwater.

--- **Glacial striation** - Point of observation is at dot. Arrow shows ice-flow direction inferred from striations on bedrock. Number is azimuth (in degrees) of flow direction.

--- **Drumlin or other glacially streamlined hill** - Symbol is parallel to glacial ice movement.

--- **Grooved till surface** - Symbol shows length and direction of narrow ridge inferred to have been carved in till by flow of glacial ice.

--- **Ice-margin position** - Line shows approximate position of ice margin during glacial retreat for major ice-margin positions.

--- **Moraine ridge** - Ridge of till, sand, and gravel deposited and/or deformed by glacial ice. Most moraines shown have been inferred from surface topography and air photo interpretation. May be underlain by Presumpscot Formation.

--- **Glaciomarine delta topset-foreset contact elevation** (in feet above modern-day sea level; from Thompson and others, 1989).

--- **Boundary of large gravel pit.**

USES OF SURFICIAL GEOLOGY MAPS

A surficial geology map shows all the loose materials such as till (commonly called lardpan), sand and gravel, or clay, which overlie solid ledge (bedrock). Bedrock outcrops and areas of abundant bedrock outcrops are shown on the map, but varieties of the bedrock are not distinguished (refer to bedrock geology map). Most of the surficial materials are deposits formed by glacial and deglacial processes during the last stage of continental glaciation, which began about 25,000 years ago. The remainder of the surficial deposits are the products of postglacial geologic processes, such as river floodplains, or are attributed to human activity, such as fill or other land-modifying features.

The map shows the areal distribution of the different types of glacial features, deposits, and landforms as described in the map explanation. Features such as striations and moraines can be used to reconstruct the movement and position of the glacier and its margin, especially as the ice sheet melted. Other ancient features include shorelines and deposits of glacial lakes or the glacial sea, now long gone from the state. This glacial geologic history of the quadrangle is useful to the larger understanding of past earth climate, and how our region of the world underwent recent geologically significant climatic and environmental changes. We may then be able to use this knowledge in anticipation of future similar changes for long-term planning efforts, such as coastal development or waste disposal.

Surficial geology maps are often best used in conjunction with related maps such as surficial materials maps or significant sand and gravel aquifer maps for anyone wanting to know what lies beneath the land surface. For example, these maps may aid in the search for water supplies, or economically important deposits such as sand and gravel for aggregate or clay for bricks or pottery. Environmental issues such as the location of a suitable landfill site or the possible spread of contaminants are directly related to surficial geology. Construction projects such as locating new roads, excavating foundations, or siting new homes may be better planned with a good knowledge of the surficial geology of the site. Refer to the list of related publications below.

OTHER SOURCES OF INFORMATION

- Hildreth, C. T., 2003. Surficial geology of the Purgatory 7.5-minute quadrangle, Androscoggin, Kennebec, and Sagadahoc Counties, Maine: Maine Geological Survey, Open-File Report 03-61, 5 p.
- Locke, D. B., and Hildreth, C. T., 2004. Surficial materials of the Purgatory quadrangle, Maine: Maine Geological Survey, Open-File Map 04-43.
- Neil, C. D., 1999. Significant sand and gravel aquifers of the Purgatory quadrangle, Maine: Maine Geological Survey, Open-File Map 99-35.
- Thompson, W. B., 1979. Surficial geology handbook for coastal Maine: Maine Geological Survey, 68 p. (out of print).

Purgatory Quadrangle, Maine

Surficial geologic mapping by

Carol T. Hildreth

Digital cartography by:

Susan S. Tolman
John B. Poisson


Robert G. Marvinyne

State Geologist

Cartographic design and editing by:

Robert D. Tucker

Funding for the preparation of this map was provided in part by the U. S. Geological Survey STATEMAP Program, Cooperative Agreement No. 02HQAG0032.



Maine Geological Survey

Address: 22 State House Station, Augusta, Maine 04333
Telephone: 207-287-2801 E-mail: mgs@maine.gov
Home page: <http://www.maine.gov/doc/nrm/nrm.htm>

Open-File No. 05-46 2005

This map supersedes Open-File Map 03-60.
For additional information, see Open-File Report 03-61.

SURFICIAL GEOLOGY OF MAINE

Continental glaciers like the ice sheet now covering Antarctica probably extended across Maine several times during the Pleistocene Epoch, between about 1.5 million and 10,000 years ago. The slow-moving ice superficially changed the landscape as it scraped over mountains and valleys (Figure 1), eroding and transporting boulders and other rock debris for miles (Figure 2). The sediments that cover much of Maine are largely the product of glaciation. Glacial ice deposited some of these materials, while others washed into the sea or accumulated in meltwater streams and lakes as the ice receded. Earlier stream patterns were disrupted, creating hundreds of ponds and lakes across the state. The map at left shows the pattern of glacial sediments in the Purgatory quadrangle.

The most recent "Ice Age" in Maine began about 25,000 years ago, when an ice sheet spread southward over New England (Stone and Borns, 1986). During its peak, the ice was several thousand feet thick and covered the highest mountains in the state. The weight of this huge glacier actually caused the land surface to sink hundreds of feet. Rock debris frozen into the base of the glacier abraded the bedrock surface over which the ice flowed. The grooves and fine scratches (striations) resulting from this scraping process are often seen on freshly exposed bedrock, and they are important indicators of the direction of ice movement (Figure 3). Erosion and sediment deposition by the ice sheet combined to give a streamlined shape to many hills, with their long dimension parallel to the direction of ice flow. Some of these hills (drumlins) are composed of dense glacial sediment (till) plastered under great pressure beneath the ice.

A warming climate forced the ice sheet to start receding as early as 21,000 years ago, soon after it reached its southernmost position on Long Island (Sarkin, 1986). The edge of the glacier withdrew from the continental shelf east of Long Island and reached the present position of the Maine coast by 13,800 years ago (Dorion, 1993). Even though the weight of the ice was removed from the land surface, the Earth's crust did not immediately spring back to its normal level. As a result, the sea flooded much of southern Maine as the glacier retreated to the northwest. Ocean waters extended far up the Kennebec and Penobscot valleys, reaching present elevations of up to 420 feet in the central part of the state.

Great quantities of sediment washed out of the melting ice and into the sea, which was in contact with the receding glacier margin. Sand and gravel accumulated as deltas (Figure 4) and submarine fans where streams discharged along the ice front, while the finer silt and clay dispersed across the ocean floor. The shells of clams, mussels, and other invertebrates are found in the glaciomarine clay that blankets lowland areas of southern Maine. Age dates on these fossils tell us that ocean waters covered parts of Maine until about 11,000 years ago, when the land surface rebounded as the weight of the ice sheet was removed.

Meltwater streams deposited sand and gravel in tunnels within the ice. These deposits remained as ridges (eskers) when the surrounding ice disappeared (Figure 5). Maine's esker systems can be traced for up to 100 miles, and are among the longest in the country.

Other sand and gravel deposits formed as mounds (barnes) and terraces adjacent to melting ice, or as outwash in valleys in front of the glacier. Many of these water-laid deposits are well layered, in contrast to the chaotic mixture of boulders and sediment of all sizes (till) that was released from dirty ice without subsequent reworking. Ridges consisting of till or washed sediments (moraines) were constructed along the ice margin in places where the glacier was still actively flowing and conveying rock debris to its terminus. Moraine ridges are abundant in the zone of former marine submergence, where they are useful indicators of the pattern of ice retreat (Figure 6).

The last remnants of glacial ice probably were gone from Maine by 10,000 years ago. Large sand dunes accumulated in late-glacial time as winds picked up outwash sand and blew it onto the east sides of river valleys, such as the Androscoggin and Saco valleys (Figure 7). The modern stream network became established soon after deglaciation, and organic deposits began to form in peat bogs, marshes, and swamps. Tundra vegetation bordering the ice sheet was replaced by changing forest communities as the climate warmed (Davis and Jacobson, 1985). Geologic processes are by no means dormant today, however, since rivers and wave action modify the land (Figure 8), and worldwide sea level is gradually rising against Maine's coast.

References Cited

- Davis, R. B., and Jacobson, G. L., Jr., 1985. Late-glacial and early Holocene landscapes in northern New England and adjacent areas of Canada. *Quaternary Research*, v. 23, p. 341-368.
- Dorion, C. C., 1993. A chronology of deglaciation and accompanying marine transgression in Maine. *Geological Society of America, Abstracts with Programs*, v. 25, no. 2, p. 12.
- Sarkin, L., 1986. Pleistocene stratigraphy of Long Island, New York, in Caldwell, D. W. (editor), *The Wisconsin stage of the first geological district, eastern New York*. New York State Museum, Bull. 455, p. 6-21.
- Stone, B. D., and Borns, H. W., Jr., 1986. Pleistocene glacial and interglacial stratigraphy of New England, Long Island, and adjacent Georges Bank and Gulf of Maine, in Shrivava, V., Bowen, D. Q., and Richmond, G. M. (editors), *Quaternary glaciations in the northern hemisphere*. *Quaternary Science Reviews*, v. 5, p. 39-52.



Figure 1: "The Bubbles" and Jordan Pond in Acadia National Park. These hills and valleys were sculpted by glacial erosion. The pond was dammed behind a moraine ridge during retreat of the ice sheet.



Figure 2: Dagget's Rock in Phillips. This is the largest known glacially transported boulder in Maine. It is about 100 feet long and estimated to weigh 8,000 tons.



Figure 3: Granite ledge in Westbrook, showing polished and grooved surface resulting from glacial abrasion. The grooves and shape of the ledge indicate ice flow toward the southeast.



Figure 4: Glaciomarine delta in Franklin, formed by sand and gravel washing into the ocean from the glacier margin. The flat delta top marks approximate former sea level. Kettle hole in foreground was left by melting of ice.




Figure 5: Esker cutting across Kearsar Five Ponds, Watford. The ridge consists of sand and gravel deposited by meltwater flowing in a tunnel beneath glacial ice.



Figure 6: Aerial view of moraine ridges in blueberry field, Sedgwick (note dirt road in upper right for scale). Each bouldery ridge marks a position of the retreating glacier margin. The ice receded from right to left.



Figure 7: Sand dune in Wayne. This and other "deserts" in Maine formed as windblown sand in late-glacial time blew out of valleys, often depositing it as dune fields on hillside downwind. Some dunes were reactivated in historical time when grazing animals stripped the vegetation cover.



Figure 8: Songo River delta and Songo Beach, Sebago Lake State Park, Naples. These deposits are typical of geological features formed in Maine since the Ice Age.